

Calculations of the Imaginary Part of Gauge Boson's Wave-function Renormalization Constants and the Modification of hep-ph/0301090

Yong Zhou^a

^a*Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China*
(Jun 5, 2003)

We have calculated the imaginary part of W and Z gauge boson's wave-function renormalization constants (wrc.) in the complex-pole mass renormalization scheme and made atonement for the paper hep-ph/0301090 where the imaginary part of gauge boson's wrc. has been ignored in the second renormalization scheme in section 4. But the conclusion of this section in hep-ph/0301090 doesn't change.

11.10.Gh, 12.15.Lk

In the complex-pole mass renormalization scheme [1], the gauge boson's wave-function renormalization constant (wrc.) should be complex number. If the gauge boson's self-energy function is written as [2]

$$\begin{array}{c} V, \mu \\ \sim \sim \sim \\ k \end{array} \bullet \begin{array}{c} V, \nu \\ \sim \sim \sim \end{array} = -i(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2})\Sigma_{VV}^T(k^2) - i\frac{k_\mu k_\nu}{k^2}\Sigma_{VV}^L(k^2) \quad (1)$$

then in the complex-pole mass renormalization scheme the form of the gauge boson's propagator in resonant region should be

$$\frac{1}{p^2 - M_0^2 + \Sigma_{VV}^T(p^2)} \sim \frac{1}{m^2 - M_0^2 + \Sigma_{VV}^T(m^2) + (p^2 - m^2)(1 + \Sigma_{VV}^{T'}(m^2))} = \frac{(1 + \Sigma_{VV}^{T'}(m^2))^{-1}}{p^2 - m^2 + B} \quad (2)$$

with $B = (m^2 - M_0^2 + \Sigma_{VV}^T(m^2))/(1 + \Sigma_{VV}^{T'}(m^2))$ Thus we obtain the gauge boson's wrc.

$$Z = (1 + \Sigma_{VV}^{T'}(m^2))^{-1} \xrightarrow{\text{one-loop}} 1 - \frac{\partial}{\partial p^2} \Sigma_{VV}^T(m^2) \quad (3)$$

In figure 1 we show the W gauge boson's self-energy diagrams which have contribution to the ξ_W -dependent imaginary part of W gauge boson's wrc..




FIG. 1. W boson's self-energy diagrams which have contribution to the ξ_W -dependent imaginary part of W boson's wrc..

Based on Eq.(1),(3) and the *cutting rules* [3], we have

$$Im[Z_W] = -\frac{e^2(1 - \xi_W)(\xi_W^2 - 2\xi_W - 11)\theta[1 - \xi_W]}{96\pi} \quad (4)$$

In figure 2 we show the Z gauge boson's self-energy diagrams which have contribution to the ξ_W -dependent imaginary part of Z gauge boson's wrc..




FIG. 2. Z boson's self-energy diagrams which have contribution to the ξ_W -dependent imaginary part of Z boson's wrc..

Based on Eq.(1),(3) and *cutting rules*, we also have

$$Im[Z_Z] = -\frac{1}{96\pi c_W^2 s_W^2} e^2 (1 - 4c_W^2 \xi_W)^{3/2} \theta[1 - 4c_W^2 \xi_W] + \frac{1}{48\pi c_W^2} e^2 ((\xi_W - 1)^2 c_W^4 - 2(\xi_W - 5)c_W^2 + 1) \sqrt{(\xi_W - 1)^2 c_W^4 - 2(\xi_W + 1)c_W^2 + 1} \theta[1/c_W - \sqrt{\xi_W} - 1] \quad (5)$$

Such imaginary part of gauge boson's wrc. will contribute to the amplitudes of physical processes, e.g. $W^- \rightarrow e_i \bar{\nu}_i$ and $Z \rightarrow d_i \bar{d}_i$. These two processes have been calculated in section 4 of hep-ph/0301090, but the imaginary part of gauge boson's wrc. has been ignored in the complex-pole mass renormalization scheme. So Eq.(57)-(60) of hep-ph/0301090 need to be modified. Eq.(58) and (60) should be gotten rid of. Eq.(57) about the imaginary part of the amplitude of $W^- \rightarrow e_i \bar{\nu}_i$ in complex-pole mass renormalization scheme should be changed as follows

$$Im[T'_1] = A_L \frac{e^3 (1 - \xi_W)^3 \theta[1 - \xi_W]}{64\sqrt{2}\pi s_W} + B_L \frac{e^3 (\xi_W - 1)^3 m_{l,i} M_W \theta[1 - \xi_W]}{32\sqrt{2}\pi s_W (M_W^2 - m_{l,i}^2)} \quad (6)$$

It is also a gauge-dependent quantity. For the process $Z \rightarrow d_i \bar{d}_i$, Eq.(59) is only the result of the on-shell mass renormalization scheme. In the complex-pole scheme, the right result is

$$Im[\mathcal{M}(Z \rightarrow d_i \bar{d}_i)]_{m_{d,i} \rightarrow 0} = A_L [-\frac{1}{576\pi c_W^3 s_W} e^3 (1 - 4c_W^2 \xi_W)^{3/2} \theta[1 - 4c_W^2 \xi_W] + \frac{1}{288\pi c_W^3} e^3 s_W ((\xi_W - 1)^2 c_W^4 - 2(\xi_W - 5)c_W^2 + 1) \times \sqrt{(\xi_W - 1)^2 c_W^4 - 2(\xi_W + 1)c_W^2 + 1} \theta[1/c_W - \sqrt{\xi_W} - 1]] + A_R [-\frac{1}{576\pi c_W^3 s_W} e^3 (1 - 4c_W^2 \xi_W)^{3/2} \theta[1 - 4c_W^2 \xi_W] + \frac{1}{288\pi c_W^3} e^3 s_W ((\xi_W - 1)^2 c_W^4 - 2(\xi_W - 5)c_W^2 + 1) \times \sqrt{(\xi_W - 1)^2 c_W^4 - 2(\xi_W + 1)c_W^2 + 1} \theta[1/c_W - \sqrt{\xi_W} - 1]] \quad (7)$$

where $A_L = \bar{d}_i \not{\epsilon}_L d_i$, $A_R = \bar{d}_i \not{\epsilon}_R d_i$. It is also gauge-dependent. At one-loop level all of these gauge-dependent terms cannot contribute to the modulus square of the physical amplitude. But we haven't still known how to cancel out the contributions of these gauge-dependent terms to the modulus square of the physical amplitude at two loop level. It needs to be done in the future.

Thus the conclusion of section 4 of hep-ph/0301090 is still alive. That's: based on the present knowledge we cannot judge which scheme is the better, on-shell mass renormalization scheme or complex-pole mass renormalization scheme.

Acknowledgments

The author thanks professor Xiao-Yuan Li for the enlightened discussion with him.

- [1] A.Sirlin, Phys. Rev. Lett. **67** ,2127(1991);
B.A.Kniehl and A.Sirlin, Phys. Rev. Lett. **81** ,1373(1998);
A.Freitas, W.Hollik, W.Walter and G.Weiglein, Phys. Lett. B495 (2000) 338.
- [2] A. Denner, Fortschr. Phys. **41** (1993) 307.
- [3] R.E.Cutkosky, J. Math. Phys. **1** (1960) 429.